



BRIGHT FUTURES HANDOUT ► PARENT

15 MONTH VISIT

Here are some suggestions from Bright Futures experts that may be of value to your family.

✓ TALKING AND FEELING

- Try to give choices. Allow your child to choose between 2 good options, such as a banana or an apple, or 2 favorite books.
- Know that it is normal for your child to be anxious around new people. Be sure to comfort your child.
- Take time for yourself and your partner.
- Get support from other parents.
- Show your child how to use words.
 - Use simple, clear phrases to talk to your child.
 - Use simple words to talk about a book's pictures when reading.
 - Use words to describe your child's feelings.
 - Describe your child's gestures with words.

✓ A GOOD NIGHT'S SLEEP

- Put your child to bed at the same time every night. Early is better.
- Make the hour before bedtime loving and calm.
- Have a simple bedtime routine that includes a book.
- Try to tuck in your child when he is drowsy but still awake.
- Don't give your child a bottle in bed.
- Don't put a TV, computer, tablet, or smartphone in your child's bedroom.
- Avoid giving your child enjoyable attention if he wakes during the night. Use words to reassure and give a blanket or toy to hold for comfort.

✓ TANTRUMS AND DISCIPLINE

- Use distraction to stop tantrums when you can.
- Praise your child when she does what you ask her to do and for what she can accomplish.
- Set limits and use discipline to teach and protect your child, not to punish her.
- Limit the need to say "No!" by making your home and yard safe for play.
- Teach your child not to hit, bite, or hurt other people.
- Be a role model.

✓ HEALTHY TEETH

- Take your child for a first dental visit if you have not done so.
- Brush your child's teeth twice each day with a small smear of fluoridated toothpaste, no more than a grain of rice.
- Wean your child from the bottle.
- Brush your own teeth. Avoid sharing cups and spoons with your child. Don't clean her pacifier in your mouth.

Helpful Resources: Poison Help Line: 800-222-1222

Information About Car Safety Seats: www.safercar.gov/parents | Toll-free Auto Safety Hotline: 888-327-4236

15 MONTH VISIT—PARENT



SAFETY

- Make sure your child's car safety seat is rear facing until he reaches the highest weight or height allowed by the car safety seat's manufacturer. In most cases, this will be well past the second birthday.
- Never put your child in the front seat of a vehicle that has a passenger airbag. The back seat is the safest.
- Everyone should wear a seat belt in the car.
- Keep poisons, medicines, and lawn and cleaning supplies in locked cabinets, out of your child's sight and reach.
- Put the Poison Help number into all phones, including cell phones. Call if you are worried your child has swallowed something harmful. Don't make your child vomit.
- Place gates at the top and bottom of stairs. Install operable window guards on windows at the second story and higher. Keep furniture away from windows.
- Turn pan handles toward the back of the stove.
- Don't leave hot liquids on tables with tablecloths that your child might pull down.
- Have working smoke and carbon monoxide alarms on every floor. Test them every month and change the batteries every year. Make a family escape plan in case of fire in your home.

WHAT TO EXPECT AT YOUR CHILD'S 18 MONTH VISIT

We will talk about

- Handling stranger anxiety, setting limits, and knowing when to start toilet training
- Supporting your child's speech and ability to communicate
- Talking, reading, and using tablets or smartphones with your child
- Eating healthy
- Keeping your child safe at home, outside, and in the car

Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 4th Edition

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit*, 2nd Edition.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.

DTaP (Diphtheria, Tetanus, Pertussis) Vaccine: *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

DTaP vaccine can prevent **diphtheria, tetanus, and pertussis**.

Diphtheria and pertussis spread from person to person. Tetanus enters the body through cuts or wounds.

- **DIPHTHERIA (D)** can lead to difficulty breathing, heart failure, paralysis, or death.
- **TETANUS (T)** causes painful stiffening of the muscles. Tetanus can lead to serious health problems, including being unable to open the mouth, having trouble swallowing and breathing, or death.
- **PERTUSSIS (aP)**, also known as “whooping cough,” can cause uncontrollable, violent coughing that makes it hard to breathe, eat, or drink. Pertussis can be extremely serious especially in babies and young children, causing pneumonia, convulsions, brain damage, or death. In teens and adults, it can cause weight loss, loss of bladder control, passing out, and rib fractures from severe coughing.

2. DTaP vaccine

DTaP is only for children younger than 7 years old. Different vaccines against tetanus, diphtheria, and pertussis (Tdap and Td) are available for older children, adolescents, and adults.

It is recommended that children receive 5 doses of DTaP, usually at the following ages:

- 2 months
- 4 months
- 6 months
- 15–18 months
- 4–6 years

DTaP may be given as a stand-alone vaccine, or as part of a combination vaccine (a type of vaccine that combines more than one vaccine together into one shot).

DTaP may be given at the same time as other vaccines.

3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of any vaccine that protects against tetanus, diphtheria, or pertussis**, or has any **severe, life-threatening allergies**
- Has had a **coma, decreased level of consciousness, or prolonged seizures within 7 days after a previous dose of any pertussis vaccine (DTP or DTaP)**
- Has **seizures or another nervous system problem**
- Has ever had **Guillain-Barré Syndrome** (also called “GBS”)
- Has had **severe pain or swelling after a previous dose of any vaccine that protects against tetanus or diphtheria**

In some cases, your child’s health care provider may decide to postpone DTaP vaccination until a future visit.

Children with minor illnesses, such as a cold, may be vaccinated. Children who are moderately or severely ill should usually wait until they recover before getting DTaP vaccine.

Your child’s health care provider can give you more information.



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

4. Risks of a vaccine reaction

- Soreness or swelling where the shot was given, fever, fussiness, feeling tired, loss of appetite, and vomiting sometimes happen after DTaP vaccination.
- More serious reactions, such as seizures, non-stop crying for 3 hours or more, or high fever (over 105°F) after DTaP vaccination happen much less often. Rarely, vaccination is followed by swelling of the entire arm or leg, especially in older children when they receive their fourth or fifth dose.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636** (**1-800-CDC-INFO**) or
 - Visit CDC's website at www.cdc.gov/vaccines.



Haemophilus influenzae type b (Hib) Vaccine: What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

Hib vaccine can prevent *Haemophilus influenzae* type b (Hib) disease.

Haemophilus influenzae type b can cause many different kinds of infections. These infections usually affect children under 5 years of age but can also affect adults with certain medical conditions. Hib bacteria can cause mild illness, such as ear infections or bronchitis, or they can cause severe illness, such as infections of the blood. Severe Hib infection, also called “invasive Hib disease,” requires treatment in a hospital and can sometimes result in death.

Before Hib vaccine, Hib disease was the leading cause of bacterial meningitis among children under 5 years old in the United States. Meningitis is an infection of the lining of the brain and spinal cord. It can lead to brain damage and deafness.

Hib infection can also cause:

- Pneumonia
- Severe swelling in the throat, making it hard to breathe
- Infections of the blood, joints, bones, and covering of the heart
- Death

2. Hib vaccine

Hib vaccine is usually given in 3 or 4 doses (depending on brand).

Infants will usually get their first dose of Hib vaccine at 2 months of age and will usually complete the series at 12–15 months of age.

Children between 12 months and 5 years of age who have not previously been completely vaccinated against Hib may need 1 or more doses of Hib vaccine.

Children over 5 years old and adults usually do not receive Hib vaccine, but it might be recommended for older children or adults whose spleen is damaged or has been removed, including people with sickle cell disease, before surgery to remove the spleen, or following a bone marrow transplant. Hib vaccine may also be recommended for people 5 through 18 years old with HIV.

Hib vaccine may be given as a stand-alone vaccine, or as part of a combination vaccine (a type of vaccine that combines more than one vaccine together into one shot).

Hib vaccine may be given at the same time as other vaccines.

3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of Hib vaccine**, or has any **severe, life-threatening allergies**

In some cases, your health care provider may decide to postpone Hib vaccination until a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting Hib vaccine.

Your health care provider can give you more information.



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

4. Risks of a vaccine reaction

- Redness, warmth, and swelling where the shot is given and fever can happen after Hib vaccination.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636** (1-800-CDC-INFO) or
 - Visit CDC's website at www.cdc.gov/vaccines.



Pediatric Dental Care

Now is the time to schedule your child's dental visit!

The AAP now recommends that all children regardless of age should use fluoride toothpaste as soon as they have teeth.

(A grain size amount for children under 3 and pea size for 3 and older)

Routine dental visits are recommended at 12 months of age then every 6 mo.



If you would like to schedule your pediatric dental visit, please call:

721-8939 – Follow the prompts for the Dental Department

**Or stop by our dental clinic here at our CHCBH site
(350 Pine Street)**

This Page Left Intentionally Blank

After the Shots...

Your child may need extra love and care after getting vaccinated. Some vaccinations that protect children from serious diseases also can cause discomfort for a while. Here are answers to questions many parents have after their children have been vaccinated. If this sheet doesn't answer your questions, call your healthcare provider.

**Vaccinations may hurt a little ...
but disease can hurt a lot!**

Call your healthcare provider right away if you answer "yes" to any of the following questions:

- ☐ Does your child have a temperature that your healthcare provider has told you to be concerned about?
- ☐ Is your child pale or limp?
- ☐ Has your child been crying for more than 3 hours and just won't quit?
- ☐ Is your child's body shaking, twitching, or jerking?
- ☐ Is your child very noticeably less active or responsive?

► Please see page 2 for information on the proper amount of medicine to give your child to reduce pain or fever.

What to do if your child has discomfort

I think my child has a fever. What should I do?

Check your child's temperature to find out if there is a fever. An easy way to do this is by taking a temperature in the armpit using an electronic thermometer (or by using the method of temperature-taking your healthcare provider recommends). If your child has a temperature that your healthcare provider has told you to be concerned about or if you have questions, call your healthcare provider.

Here are some things you can do to help reduce fever:

- Give your child plenty to drink.
- Dress your child lightly. Do not cover or wrap your child tightly.
- Give your child a fever- or pain-reducing medicine such as acetaminophen (e.g., Tylenol) or ibuprofen (e.g., Advil, Motrin). The dose you give your child should be based on your child's weight and your healthcare provider's instructions. See the dose chart on page 2. *Do not give aspirin.* Recheck your child's temperature after 1 hour. Call your healthcare provider if you have questions.

My child has been fussy since getting vaccinated. What should I do?

After vaccination, children may be fussy because of pain or fever. To reduce discomfort, you may want to give your child a medicine such as acetaminophen or ibuprofen. See the dose chart on page 2. *Do not give aspirin.* If your child is fussy for more than 24 hours, call your healthcare provider.

My child's leg or arm is swollen, hot, and red. What should I do?

- Apply a clean, cool, wet washcloth over the sore area for comfort.
- For pain, give a medicine such as acetaminophen or ibuprofen. See the dose chart on page 2. *Do not give aspirin.*
- If the redness or tenderness increases after 24 hours, call your healthcare provider.

My child seems really sick. Should I call my healthcare provider?

If you are worried at all about how your child looks or feels, call your healthcare provider!

HEALTHCARE PROVIDER: PLEASE FILL IN THE INFORMATION BELOW.

If your child's temperature is 104 °F or 40 °C or higher,
or if you have questions, call your healthcare provider.

Healthcare provider phone number 605-721-8939

Medicines and Doses to Reduce Pain and Fever

Choose the proper medicine, and measure the dose accurately.

1. Ask your healthcare provider or pharmacist which medicine is best for your child.
2. Give the dose based on your child's weight. If you don't know your child's weight, give the dose based on your child's age. Do not give more medicine than is recommended.
3. If you have questions about dosage amounts or any other concerns, call your healthcare provider.
4. **Always use a proper measuring device when giving acetaminophen liquid (e.g., Tylenol) or ibuprofen liquid (e.g., Advil, Motrin):**
 - Use the device enclosed in the package.
 - If you misplace the device, consult your healthcare provider or pharmacist for advice.

■ **Meal-time spoons are not accurate measures.** Never use a meal-time spoon for giving medication.

Take these two steps to avoid causing a serious medication overdose in your child.

1. Don't give your child a larger amount of acetaminophen (e.g., Tylenol) or ibuprofen (e.g., Motrin, Advil) than is shown in the table below. Too much of any of these medicines can be extremely dangerous.
2. When you give your child acetaminophen or ibuprofen, don't also give them over-the-counter cough or cold medicines. This can cause a medication overdose because cough and cold medicines often contain acetaminophen or ibuprofen. In fact, to be safe, don't ever give over-the-counter cough and cold medicines to your child unless you talk to your child's healthcare provider first.

ACETAMINOPHEN (Tylenol or another brand): How much to give?

Give every 4 to 6 hours, as needed, no more than 5 times in 24 hours (unless directed to do otherwise by your healthcare provider).

Child's weight	Child's age	Infants' or children's liquid 160 mg in each 5 mL	Children's chewables – current product 160 mg in each tablet	Infants' drops 80 mg in each 0.8 mL	Children's chewables 80 mg in each 0.8 mL
6–11 lbs (2.7–5 kg)	0–3 mos	Advised dose* _____		OLD PRODUCT	OLD PRODUCT
12–17 lbs (5.5–7.7 kg)	4–11 mos	2.5 mL		Throw away this product. It is out of date and should not be used.	Throw away this product. It is out of date and should not be used.
18–23 lbs (8.2–10.5 kg)	12–23 mos	3.75 mL			
24–35 lbs (10.9–15.9 kg)	2–3 yrs	5 mL	1 tablet		
36–47 lbs (16.4–21.4 kg)	4–5 yrs	7.5 mL	1½ tablets		
48–59 lbs (21.8–26.8 kg)	6–8 yrs	10 mL	2 tablets		
60–71 lbs (27.3–32.3 kg)	9–10 yrs	12.5 mL	2½ tablets		
72–95 lbs (32.7–43.2 kg)	11 yrs	15 mL	3 tablets		

IBUPROFEN (Advil, Motrin, or another brand): How much to give?

Give every 6 to 8 hours, as needed, no more than 4 times in 24 hours (unless directed to do otherwise by your healthcare provider).

Child's weight	Child's age	Infants' drops 50 mg in each 1.25 mL 	Children's liquid 100 mg in each 5 mL 	Children's chewables or junior tablets 100 mg in each tablet	Children's chewables 50 mg in each tablet
less than 11 lbs (5 kg)	0–5 mos				OLD PRODUCT
12–17 lbs (5.5–7.7 kg)	6–11 mos	1.25 mL	Advised dose* 2.5 mL		Throw away this product. It is out of date and should not be used.
18–23 lbs (8.2–10.5 kg)	12–23 mos	1.875 mL	Advised dose* 3.75 mL		
24–35 lbs (10.9–15.9 kg)	2–3 yrs		5 mL	1 tablet	
36–47 lbs (16.4–21.4 kg)	4–5 yrs		7.5 mL	1½ tablets	
48–59 lbs (21.8–26.8 kg)	6–8 yrs		10 mL	2 tablets	
60–71 lbs (27.3–32.3 kg)	9–10 yrs		12.5 mL	2½ tablets	
72–95 lbs (32.7–43.2 kg)	11 yrs		15 mL	3 tablets	

* **HEALTHCARE PROVIDER:** Please fill in the advised dose.

Immunization Action Coalition • www.immunize.org/catg.d/p4015.pdf • Item #P4015 (2/19)